

Full Length Research Paper

Effects of underground water on soil salinity and dates production in Kebili oases area (Tunisia): The case of El Bahaier oasis

¹Habib Ben Hassine, ¹Achwak Ben Slimane, ²Mohamed Mlawah, ¹Lassaad Albouchi and ²Ammar Gandouzi

¹University of Carthage, Graduate School of Agriculture of Mograne, 1121 Mograne, Tunisia, ²Service of soils, CRDA Kebili, Tunisia.

Accepted 17 July, 2013

The risks of increase of the level of the salted underground water and its salinization effect on the soils in oasis creations are inescapable because of their altitude which is very close to that of the next salted depressions. To control this phenomenon, some monitoring points called piezometers were installed in these perimeters, among them the one of El Bahaier located at the north of Kébili city in Tunisia and being the subject of the follow-up of the current study. Two measurement operations of the level of the underground table and its salinity were carried out in the functional piezometers of this oasis in March and April 2011; in parallel, two series of soils samples were taken around these same piezometers on a depth of 1 meter. The determination of the salinity of these soils samples showed that the most waterlogged zones are those which are salinized explaining as well as the increase of the underground water level is a principal cause of the soils salinization in these zones known by their high potential evapotranspiration and very weak rainfall. This observation is confirmed by the correlation tests between various couples of parameters, where the correlation coefficient r was generally high and significant. Moreover, the effect of the waterlogging and the soils salinity on the output of the principal culture practiced in this perimeter, the date palm (*Phoenix dactylifera*), was highly significant. The improvement, the clearing out and the cleaning of the drains are necessary to push the level of in-depth subsoil water more and to relieve the trees of its negative effect on their production. However, an optimal level of this water table located at 2 meters of depth would be more beneficial on the yield.

Keywords: Groundwater, soil, oasis, salinization, waterlogging, Tunisia.

INTRODUCTION

Salinization of irrigated soils in arid areas, which can be described as secondary salinization because it occurs due to irrigation activities, may have a double origin: (i) irrigation waters which, despite their varied salt concentrations, deposit at long-term significant amounts of salts in the growth zone of roots of cultivated plants, (ii) underground water often salty, approaching the soil

surface after receipt of excess irrigation water for salts leaching, communicate to the surface horizons of soil salts dissolved therein. This second category of salinization is usually the most dangerous due to high water concentrations of these layers and also the capillary active in environments with high evaporation. FAO (2006) estimated that the world loses an average of 3 hectares of farmland per minute by salinization and financial losses caused by the degradation factor amount to 250 \$/ha. Perimeters which are subject to regimes of salt shallow aquifers are located

*Corresponding author E-mail: habib.benhassine10@gmail.com

either in low altitude depressions characterized by deficient internal drainage or around salted areas or near coasts (Ben Hassine, 2005). The second case is that of continental oasis of Jerid and Nefzaoua which are located around the big Chott Jerid and Gharsasalt lakes, situations where the difference in elevation between the saline lake and the oasis are low causing difficulties for groundwater discharge (Ben Aissaand al., 2004; Marletand al., 2007). Almost stagnant waters of the water table are a constant source of salts to the soil in which the solutions become more concentrated and the osmotic pressures are increasing causing more difficult conditions of absorption of moisture and nutrients. The date palm culture which adapts to conditions of high summer temperatures and high salinities is the main species in these oases, often accompanied by two shrub and herbaceous (vegetable) strata (Job and Marai, 1990). It assumes important roles in the social, economic and ecological aspects (Kadri and Van Ranst, 2002; Belloumi and Matoussi, 2007). It participates in the economic activities of the country (Tunisia) through the production of 102000 tons of dates (Van Ranst and Kadri, 2002). Its safeguarding and protection against degradation factors and in particular against salinization and waterlogging of soils becomes a necessity. The first steps in this protection are installation of drainage systems for evacuating away from the oasis, excess water and salts (Côte, 1998), the proper management of irrigation water (Legros, 2009; Marlet, 2004; Marletand et al., 2007) and especially monitoring and periodic controls of the evolution of the underground water and soil salinity (Hatiraand et al., 2005).

Programs to monitor the evolution of soil salinity and underground water levels are set up in almost all oases subject to the effects of waterlogging and salinization. This monitoring is carried out by using piezometers, which are distributed in a way to cover well the perimeter controlled in terms of its surface and its waterlogged area. The current study retains one of these controlled oases, which is El Bahaier, located at the north of Kebili city in the Nefzaoua region (South of Tunisia). It is situated in the basin flow area of Chott El Fejaj. The measurements carried out in spring 2011 had the objective to assess the impact of the underground water level and salinity on soil salinity and also on the date palms production.

MATERIALS AND METHODS

Location and environmental conditions of the study area

The oasis of El Bahaier covers an area of 40ha at 31km north-east of Kebili city in South of Tunisia (figure 1). It takes place between the south Tebaga relief and the north salty depression of Chott El Fejaj. Its location at an

altitude close to that of the saline lake, makes that gravity underground water flows are quite slow, which favors the stagnation of excess water and the creation of waterlogging conditions. This oasis is planted with three vegetative strata: date palms (*Phoenix dactilifera*) aged ten years, whose density varies between 93 and 100 plants/ha, fruit trees like olive, pomegranate, fig, grapevine and yearly herbaceous crops consisting of forage and vegetable cultures. Annual rainfall of the area does not exceed 100 mm and annual potential evapotranspiration measured by the Thornthwaite formula is nearly 1300 mm (Kadri and Van Ranst, 2002; Marletand al, 2007).

This oasis is irrigated from a borehole capturing the deep aquifer of the intercalary Continental, whose concentration of salts is close to 3 g/l. Excess irrigation water caused the rise of the topped water table level. The installation of an open drainage system helped to keep this shallow groundwater in a steady state. To control the evolution of the level, 12 piezometers were located in a fairly homogeneous throughout the perimeter (figure 2). A piezometer is a cylindrical tube of metal or PVC, 8 cm in diameter and 3 m in length, which is drive down into the soil to 2,5 m depth. It is strainer on the bottom half so as to allow the penetration of water from the underground water for measuring level or sampling.

The soils of the oasis

The soil covering the oasis land is gypsiferous and encrusted alluvial deposit, salty in the deep horizons. Salinity increased after starting irrigation in the perimeter. When the crust is superficial, it is a gypsum-limestone type. The surface horizon of 30 cm thick and sandy-loam texture is covering a gypsiferous sandy and encrusted material.

This soil can be classified as "calcimagnésique gypseux" (CPCS, 1967) or Haplic GYPSOSOL, limestone with gypsic horizon in depth (AFES, 2008). To soil taxonomy, we can classify it among aridisols: gypsiorthids (SMSS, 1985).

Field work

Two measurement campaigns of the water level and salinity were carried out in eight functional piezometers in March and April 2011, four piezometers were indeed damaged when passing gear tillage. Level was measured using an electric probe and the sample was taken using a small container attached to a rope. Measuring salinity was performed directly on the field using a portable conductivity meter.

Meanwhile, two series of soil samples were carried out during the same periods around the eight functional piezometers. Sampling was carried out with auger every 20cm to a depth of 1 meter. In total, 80 soil samples were collected during two campaigns. The measurement




Figure 1. Location of Tunisia and the studied region in North Africa (wpmmap.org).




Figure 2. Land plot plan of El Bahaier oasis and location of the piezometers.

of soil electrical conductivity (EC) was performed on a saturated paste extract. The economic aspect was associated with the technical study through a survey on the average yield of a date

palm tree during the precedent crop, in parcels which were conducted level and salinity measurements of the groundwater and also the salinity of the different soil horizons.

Table 1. Averages of the two measures of the studied parameters, carried out in March and April 2011.

Piezometers numbers	Level of the water cm	EC of the water mS/cm	Electric conductivity (EC) of saturated paste extract (mS/cm)					Average EC*	Yield kg/tree**
			0-20cm	20-40cm	40-60cm	60-80cm	80-100 cm		
1	119,5	8,31	11,77	9,82	8,48	8,08	8,20	10,42	30
3	113,5	5,55	8,25	5,90	6,58	5,46	5,27	7,09	30
4	117,5	7,57	5,47	5,59	5,92	5,99	5,06	5,56	40
5	130	4,53	5,11	4,90	4,88	4,76	4,17	4,95	55
6	144	5,35	4,61	4,58	5,07	4,37	3,59	4,58	60
7	234,5	5,70	3,95	3,22	3,57	3,98	3,51	3,69	60
8	251	3,18	3,82	3,37	3,69	3,89	3,71	3,69	65
11	237,5	3,61	6,13	4,95	4,49	4,04	3,80	5,35	60
Average	168,44	5,47	6,14	5,29	5,33	5,07	4,66	5,66	50,00
$\sigma/n^{0,5}$	60,98	1,78	2,68	2,06	1,63	1,43	1,58	2,21	14,39
CV %	36,21	32,49	43,63	38,95	30,56	28,20	33,86	39,02	28,78

* Average EC = $(2 \cdot EC_{0-20} + EC_{20-40} + 0,5 \cdot EC_{40-60} + 0,25 \cdot EC_{60-80} + 0,25 \cdot EC_{80-100})/4$ (CRUESI, 1970)

** Yield of crop 2010 obtained by a survey with the farmers.

Statistical calculations

Data from the two measurements in March and April 2011 were expressed by their average. Then, the variation between the values of different points was illustrated by calculating the average of the standard deviation and coefficient of variation. The study of the impact of various parameters measured on each other was performed by calculating the correlation coefficients for each pair of values of the parameters taken pairwise.

RESULTS

Level and salinity of the underground water

The water level varied between 113,5 cm and 251cm. The average of all points was 168,44 cm with a standard deviation of 60,98 cm and a coefficient of variation of 36,21% (table 1). Points where the level seems closer to the surface are those numbered 1, 3, 4 and to a lesser extent the point 5. The salinity of the water is high in points 1 and 4, moderate in 3, 5, 6 and 7 and quite low in points 8 and 11 (table 1). It seems quite clear that the higher the level distance from the surface, the greater the salinity decreases and vice versa. So, underground water is more concentrated in salts when approaching the surface.

Soils salinity

Points 1 and 3 are characterized by higher salinity which reaches respectively 8,25 and 11,77 mS/cm in the surface horizons. It remains high in the underlying horizons, but the saline profile is ascendant. The lowest values are observed in points 7 and 8 where saline profiles are vertical. Points 4, 5, 6 and 11 are

intermediate salinity, which is near to 5mS/cm, their profiles are generally vertical saline, except for point 11 where the salinity of the surface horizon is significantly higher than those of other layers (table 1). The lateral and vertical distribution of salts reveals that there is, a priori relationship between the underground water level and soil salinity and over the water approaches the surface, the more saline profile tends to become ascendant. The highest value of the coefficient of variation is observed in the surface horizon (0-20cm) followed by the deep (80-100cm), the lowest variation corresponds to the horizon 60-80cm with a value of 28,20 %.

Salinity of the oasis soils, especially important around piezometers 1 and 3 where the level of underground water is closer to the surface, should act negatively on the production of date palm. Indeed, the lowest yields were obtained near of the most waterlogged and salty soils.

Effects of underground water on soil salinity

Undergroundwater acts on the soil by the rise of its waters through capillary porosity at a height which varies depending on the texture, this rise is the most active in loamy soils (Ben Hassine, 2005). The correlation coefficient (r) between the underground water level and soil salinity is always less than -0,5 explaining that there is a more or less significant correlation between the two parameters. Indeed, the greater the depth of the water decreases as salinity increases, which explains the negative value of r. The most significant correlations (threshold 0,05 at n-2 degrees of freedom) were obtained for levels 40-60cm and 60-80cm (table 2).

Salinity of the undergroundwater and salinity of soil horizons are better correlated since the electrical conductivities of the last four horizons (20-100cm) are significantly related (thresholds 0,01 and 0,05) to the


Table 2. Correlations matrix between the measured parameters.

	Water level cm	EC water mS/cm	EC(0-20) mS/cm	EC(20-40) mS/cm	EC(40-60) mS/cm	EC(60-80) mS/cm	EC(80-100) mS/cm	Average EC mS/cm	Yield kg/tree
Water level cm	1								
EC water mS/cm	-0,677	1							
EC(0-20) mS/cm	-0,540	0,622	1						
EC(20-40) mS/cm	-0,633	0,723*	0,960***	1					
EC(40-60) mS/cm	-0,775*	0,773*	0,933***	0,963***	1				
EC(60-80) mS/cm	-0,700*	0,858**	0,884**	0,953***	0,954***	1			
EC(80-100) mS/cm	-0,588	0,766*	0,941***	0,966***	0,942***	0,978***	1		
Average EC mS/cm	-0,600	0,685	0,993***	0,985***	0,962***	0,929***	0,966***	1	
Yield kg/tree	0,771*	-0,759*	-0,835**	-0,808*	-0,910**	-0,866**	-0,846**	-0,851**	1

* :significant correlation at threshold 0,05, with 6 degrees of freedom (n-2)

* * : significant correlation at threshold 0,01, with 6 degrees of freedom (n-2)

*** :significant correlation at threshold 0,001, with 6 degrees of freedom (n-2).

**Figure 3.** Correlation curve between the underground water level and yield of palm trees.

water salinity (table 2). The salt concentration of the water would have obviously an effect on the accumulation of salts in different soil depths and particularly in the 60-80cm layer. Only the surface horizon has no significant relationship with the water salinity probably because firstly of its location farther and secondly for receiving permanent irrigation water that disrupts this relationship.

The average salinity is not significantly related either to the depth of the water nor its salinity. It more or less reflects the values of the surface horizon, that are doubled in the calculating formula of the average value on each sampling point.

We must not omit to mention the very highly significant correlations between salinities of various soil horizons (table 2), which reflects the inter-communicating salinity




Figure 4. Correlation straight line between the underground water salinity and yield of palm trees.

in the profile even between the most distant horizons (0–20cm and 80–100cm).

Effect of underground water on the yield of date palm

The date palm yield increases when the water level moves away from the soil surface, to a certain limit. Indeed, the correlation curve which is polynomial and has a high coefficient of determination R^2 , increases to the level of 75,8 kg/tree and then relapse again explaining that when the water is too far from the surface, the palm tree bends in production (figure 3). The level at which yield begins to decrease is 199 cm, that is equal to approximately 2 meters. It draws the consequence that the date palm produces better when the water is at 2 meters, going up or away from the optimal level, the underground water would have a negative effect on the production of this species.

For the relationship between the salinity of the water and the dates yield, it is illustrated by the linear equation of the form $-ax + b$ (figure 4). The shape of the right means more than the water salinity is increasing the yield decreases. The relationship is significant at the 0,05 threshold. The highest concentration of salts in the underground water should create more difficult conditions of water supply and extreme toxicity for the root system of the tree.

Effect of soil salinity on the yield of date palm trees

Almost all horizons showed a significant correlation between their salinity and the yield of date palm trees (table 2). To illustrate this correlation in one figure, it has been realized between the average soil salinity and yield

(figure 5). The right is of the form $-ax + b$, and that means more soil salinity increases, yield decreases, which should reflect the difficulties of tree nutrition in solutions highly concentrated in salts.

DISCUSSION

The oasis of El Bahaier, although located outside the basin Chott El Jerid, has climate, soil and crop similarities with all those of the Nefzaoua region. Indeed, low rainfall, the very high potential evapotranspiration, soils enriched in gypsum and the proximity of saline depressions are common criteria for almost all oases of this region. Their irrigation by deep water, little or not related to groundwater, produces the rise of the water level causing both waterlogging and salinization. The installation of drainage systems has decreased at a certain depth of the underground water, but their conception as open ditches reducing their effectiveness evacuation of water excess, was behind the creation waterlogged and saline areas in these oases. This impacts on soils and crops. Indeed, the water table is not static and moves by ascendant movements by following the capillary pores of the soil. These movements are also accompanied by those of the salts dissolved in the groundwater which is more concentrated by receiving leaching waters passed through the soil (FAO, 2006).

The results obtained in this study can conclude that the effect of the underground water is evident on soil salinity both by the depth of its level or concentration of salts in its waters. The accumulation of salts in the soil is not only a degradation factor acting negatively on their physico-chemical properties but is an indication of loss of fertility




Figure 5. Correlation straight line between the average of soil salinity and yield of palm trees.

of these soils (Legros, 2009) and a weakening of the balance of the oasis system which sustainability is threatened (Ben Aissa et al., 2004).

These conditions of waterlogging and salinity affect the trees of the main crop produced in these areas characterized by specific soil and climatic conditions, disrupting their nutrition solutions with high osmotic pressures. Such conditions negatively affect the physiology of the tree which reacts in two ways (Van Ranst and Kadri, 2002):

A decrease in tree vigor by reducing the height, diminution of trunk circumference and number of green palms.

A significant reduction in the production of the trees and the quality of dates.

The yields decrease was demonstrated by the results of this study and also by those of Daddy Bouhounand et al., (2011) and Zidi (1997). Yield is indeed negatively affected by both the rise of the groundwater level, the salt concentration and high salinity of soils. Daddy Bouhounand et al., (2011) identified an additional factor which acts negatively on the yield; it is the presence of a crust of compact gypsum horizon in the root zone. The proper functioning of the drainage system and the possibility of its improvement that would lead to permanent removal of the underground water to 2 meters deep, would be a solution to both reduce the risk of capillary rise from concentrated groundwater and to not deprive the roots of the date palm trees from a beneficial source of moisture.

The case studied is one of many similar cases of soils salinization around Nefzaouachotts on initially low salted soils; this process leads to a lateral extension of waterlogging and salinization. It is a kind of human and

progressive endoreisation (Mhiriand et al., 1998) which extends the vast land of saline area in the region. More rigorous management of irrigation water and soil (Cheverry and Robert, 1998) across the entire watershed basin of Chott El Jerid is to adopt in order to minimize the spread of salinization and save resources by preserving available soils against all forms of degradation.

CONCLUSION

Problems of waterlogging and salinization of soils from saline underground water are common to all Nefzaouaoases concentrated mostly around saline depressions known locally as chotts. The oasis of El Bahaier is a representative example of these processes which are almost similar for the majority of irrigated area in the region. Altitude close enough to those of neighboring saline areas and using irrigation water pumped from deep aquifers often coupled with incomplete execution of a drainage system, make that the underground water rises progressively and will be distributed unequally in the irrigated area. The lower parts of the oasis suffers a more pronounced engorgement which leads to soil salinization. Indeed, the soil is more salty near the points where the water is closer to the surface. Statistical correlations have also confirmed this relationship because the underground water level and its salinity were almost always highly correlated with soil salinity. These parameters degradation of soil quality have impacted negatively on the main agricultural production in the region, the dates palm tree whose yield was lower in salinized and waterlogged areas. The relationship between the underground water level and the yield was

realized by a polynomial equation which showed that date palm does not provide the best production in a highly waterlogged and salty zone but prefer not survive in a dry environment with low moisture in depth. An underground water level maintained at 2 meters depth would be an optimal situation.

REFERENCES

- Association française pour l'étude du sol (AFES), 2008. Référentiel pédologique français. Editions Quæ. p.196-200
- Belloumi M et al., Matoussi MS, 2007. Impacts de la salinité sur l'efficacité technique de l'agriculture irriguée : application au cas des Oasis de Nefzaoua en Tunisie. *Économie et Prévision*, n°177 2007-1, p.77-78.
- Ben Aissa I, Bouksila F, Bahri A, Bouarfa S, Chaumont C et Hichri W, 2004. Gestion de l'eau et des sels au sein d'une oasis du Sud tunisien. Acts of seminary "Modernization of irrigated agriculture", Rabat (Morocco), 19 - 23 April 2004, 12 pages.
- Ben Hassine H, 2005. Effets de la nappe phréatique sur la salinisation des sols de cinq périmètres irrigués en Tunisie. *Étude et Gestion des Sols*, Volume 12, 4, 2005, p. 281 - 300.
- Chevry Robert M, 1998. La dégradation des sols irrigués et de la ressource en eau : une menace pour l'avenir de l'agriculture et pour l'environnement des pays au sud de la méditerranée ? *Étude et Gestion des Sols*, Volume 05, 4, 1998, p. 217 - 226.
- Commission de Pédologie et de cartographie des sols (CPCS), 1967. Classification des sols, édition 1967, INRA France, 96 pages.
- Côte M, 1998. Des oasis malades de trop d'eau ? *Revue Sécheresse*, vol 9, n° 2, 1998, p. 123-130.
- CRUESI, 1970. Research and training on irrigation with saline water. Centre de recherche sur l'utilisation des eaux saumâtres en irrigation (CRUESI), technical report, UNESCO-Paris / UNDP, 1962-1969, 256 pages.
- Daddi Bouhoun M, Marlet S, Brinis L, Saker ML, Rabier J and Côte M, 2010. A survey of the combined effects of waterlogging and salinity on fruit yield in the date palm groves of the Wargla basin, Algeria. *Revue Fruits*, 2011, vol. 66, p. 11-24.
- FAO, IPTRID, 2006. Electronic conference on salinization : extension of salinization and strategies of prevention and rehabilitation. Project CISEAU. www.agrireseau.qc.ca/agroenvironnement...Salinisation_irrigation
- Hatira A, Benmansour B, Grira M et al., Gallali T, 2005. Impact des eaux d'irrigation sur l'origine des accumulations gypseuses dans les sols de l'oasis de Metouia (Tunisie). *Étude et Gestion des Sols*, Volume 12, 1, 2005, p. 43- 54.
- Job JO et al., Marai M, 1990. Etude de la salinité des sols de l'oasis El Guettar. ES 258, Soils directorate, Ministry of agriculture, Tunisia - Centre ORSTOM, Tunis, 14pages.
- Kadri A et al., Van Ranst, 2002. Contraintes de la production oasienne et stratégies pour un développement durable. Cas des oasis de Nefzaoua (Sudtunisien). *Science et changements planétaires / Sécheresse*. Volume 13, Numéro 1, p. 5-12, march 2002, Notes méthodologiques.
- Legros JP, 2009. La salinisation des terres dans le monde. *Académie des Sciences et Lettres de Montpellier*, 2009, <http://academie.biu-montpellier.fr/>, conference n°4069, Bull. n°40, p. 257-269.
- Marlet S, 2004. Evolution des systèmes d'irrigation et gestion de la salinité des terres irriguées. Acts of seminary "Modernization of irrigated agriculture", Rabat (Morocco), 19 - 23 April 2004, 11 pages.
- Marlet S, Bouksila F, Mekki I et Ben Aissa I, 2007. Fonctionnement et salinité de la nappe de l'oasis de Fatnassa : arguments géochimiques. Acts of the third regional workshop of Sirma project, Nabeul, Tunisia, 4-7 June 2007, 14 pages.
- Mhiri A, Tarhouni J, Hachicha M et Lebdi F, 1998. Approche systémique des risques de salinisation par endoréisation anthropique. *Étude et Gestion des Sols*, Volume 05, 4, 1998, p. 257-268.
- Soil Management Support Service (SMSS), 1985. Keys to soil taxonomy, technical monograph N° 6, second printing, p. 91-106.
- Zidi C, 1997. Régime de la nappe superficielle et incidences sur la salure des sols et la production des dattes dans l'oasis de Tarfaya (Kébili). ES 297, Soils directorate, Ministry of agriculture, Tunisia. Wpmap.org. Tunisia_intro_map.jpg.