

Review

Production of bioethanol from sweet sorghum: A review

A. Almodares^{1*} and M. R. Hadi²

¹Department of Biology, University of Isfahan, Isfahan-Iran.

²Department of Biology, Sciences and Research Branch of Fars, Islamic Azad University, Iran.

Accepted 13 August, 2018

The consumption of bioethanol as biofuel may reduce greenhouse gases, gasoline imports. Also it can be replaced with lead or MTBE (Methyl tert-butyl ether) that are air and underground water pollutants, respectively. Plants are the best choice for meeting the projected bioethanol demands. For this scope, a comparative analysis of the technological options using different feedstocks should be performed. Our research and other studies indicate that sweet sorghum can be used as a feedstock for ethanol production under hot and dry climatic conditions. Because, it has higher tolerance to salt and drought comparing to sugarcane and corn that are currently used for biofuel production in the world. In addition, high carbohydrates content of sweet sorghum stalk are similar to sugarcane but its water and fertilizer requirements are much lower than sugarcane. Also, sugarcane is not a salt tolerant plant. On the other hand, high fermentable sugar content in sweet sorghum stalk makes it to be more suitable for fermentation to ethanol. Therefore, it is suggested to plant sweet sorghum for biofuel production in hot and dry countries to solve problems such as increasing the octane of gasoline and to reduce greenhouse gases and gasoline imports.

Key words: Sweet sorghum, carbohydrate, bioethanol, biofuel.

INTRODUCTION

Each year, fossil energy resource is reducing in the world. Therefore, a substitute should be found. There are many crops available for producing energy such as sweet sorghum which not only produce food (Anglani, 1998), but also energy (Reddy et al., 2005), feed (Almodares et al., 1999; Fazaeli et al., 2006) and fiber (Murray et al., 2008a,b). Sorghum can be classified as sweet, grain and forage types (Almodares et al., 2008b). Sweet sorghum like grain sorghum produces grain 3 - 7 t/ha (Almodares and Mostafafi, 2006). But the essence of sweet sorghum is not from its seed, but from its stalk, which contains high sugar content (Almodares et al., 2008c). In general, it can produce stalk 54 - 69 t/ha (Table 1) (Almodares et al., 2008c).

The sugar content in the juice of sweet sorghum varies in different varieties (Almodares et al., 1994a). The Brix range in different varieties of sweet sorghum is 14.32 -

22.85% (Table 2) (Almodares and Sepahi, 1996).

Besides having rapid growth, high sugar accumulation (Almodares and Sepahi, 1996), and biomass production potential (Almodares et al., 1994a), sweet sorghum has wider adaptability (Reddy et al., 2005). Also it is well adapted to sub-tropical and temperate regions of the world and it is water efficient. Sweet sorghum has many good characteristics such as a drought resistance (Tesso et al., 2005), waterlogging tolerance, salinity resistance (Almodares et al., 2007a; Almodares et al., 2008) and with a high yield of biomass etc.

In addition, sweet sorghum is a C4 crop with high photosynthetic efficiency. Thus development of sweet sorghum will play an important role in promoting the development of agricultural production, livestock husbandry (Fazaeli et al., 2006), energy sources (biofuel) (Nahvi et al., 1994a, b), refining sugar, paper making etc. Carbohydrates, which are present in sweet sorghum, can be nonstructural such as sugars and starch, or structural such as cellulose, hemicellulose, and pectic substances (Anglani, 1998). The chief sugars present in sorghum kernels are the monosaccharides glucose and fructose,

*Corresponding author: E-mail: aalmodares@yahoo.com. Tel.: +98-3117932794. Fax: +98-311-7932792.

Table 1. Comparison of sugarcane, sugar beet and sweet sorghum in Iran.

	Sugarcane	Sugar beet	Sweet sorghum
Crop duration	About 7 months	About 5 - 6 months	About 4 months.
Growing season	Only one season	Only one season	One season in temperate and two or three seasons in tropical area.
Soil requirement	Grows well in drain soil	Grows well in sandy loam; also tolerates alkalinity	All types of drained soil.
Water management	36000 m ³ /h	18000 m ³ /h	12000 m ³ /h
Crop management	Requires good management	Greater fertilizer requirement; requires moderate management	Little fertilizer required; less pest and disease complex; easy management.
Yield per ha	70 - 80 tons	30 - 40 tons	54 - 69 tons.
Sugar content on weight basis	10 - 12%	15 - 18%	7 - 12%.
Sugar yield	7 - 8 tons/ha	5 - 6 tons/ha	6 - 8 tons/ha.
Ethanol production directly from juice	3000 - 5000 L/ha	5000 - 6000 L/ha	3000 L/ha.
Harvesting	Mechanical harvested	Very simple; normally manual	Very simple; both manual and through mechanical harvested.

the disaccharides sucrose and maltose and the trisaccharide raffinose. According to the kind of sugar in the stalk, it can be divided into saccharin- type sweet sorghum and syrup-type sweet sorghum (Anglani, 1998). Saccharin-type sweet sorghum, which mainly contains sucrose, can be used for refining crystal sugar. Syrup-type sweet sorghum, which mainly contains glucose, can be used for producing syrup. Sugars content in sweet sorghum stalk juice mostly were sucrose and invert sugars which invert sugars are included glucose, fructose, maltose and xylose (Almodares et al., 2008c). Also, they reported that mannose, galactose and arabinose were not detected in sweet sorghum juice. Therefore, it seems that using carbohydrates in the stalk (sucrose and invert sugar) is suitable for ethanol production for biofuel because these carbohydrates are easily converted to ethanol. Although, ethanol can be produced from sweet sorghum grain (Figure 1) but it needs more process for converting its starch to glucose that later will be converted to ethanol (Jacques et al., 1999).

In addition, the produced baggass after juice extraction can be used for ethanol production (Jacques et al., 1999) or animal feed (Jafarinia et al., 2005). However, presently it is not economically feasible to produce ethanol from sweet sorghum baggass (Drapcho et al., 2008). The aim of this review is to summarize the information available on sorghum carbohydrates for biofuel production.

Sweet sorghum agronomy

Sweet sorghum cultivation and practices are simple and readily adoptable (Almodares et al., 1997b). It is a short –

day plant (Almodares et al., 2000; Rezaie et al., 2005), and most varieties require fairly high temperature (Reisi and Almodares, 2008) to make their best growth. The cereals (Tesso et al., 2005) and tolerate a wide range of soil conditions (Almodares et al., 2008e). Sorghum tolerates compacted subsoil and can stand high press wheel pressure at planting. It tolerates a pH range of 5.0 to 8.5 (Smith and Frederiksen, 2000) and some degree of salinity (Almodares et al., 2007a, 2008a, 2008c, 2008), alkalinity and poor drainage (Almodares et al., 2008e). It also will grow on heavy, deep cracking vertisols and light sands (Smith and Frederiksen, 2000). The seed of sweet sorghum should be planted deep enough to give it moisture to germinate and allow its roots to grow down through moist soil into subsoil moisture, ahead of the drying front (Almodares et al., 2008e).

Planting time (Almodares and Mostafafi, 2006) usually start when the air temperature is above 12°C (Almodares et al., 2008e). Late planting reduces the length of the growing season, yield and carbohydrate content (Almodares et al., 1994a). Also, it may cause late and troublesome harvest and may expose the crop to pests and diseases and other hazards which are dominant at the end of the crop season (Almodares et al., 2008e). Balanced fertilization can increase yield (Rego et al., 2003). Nitrogen fertilizer and its application time (Almodares et al., 1996) promotes sucrose content and growth rate in sweet sorghum (Tsialtas and Maslaris, 2005). Application of adequate amounts of K fertilizer increase yield responses than increasing levels of nitrogen fertilizer alone (Pholsen and Sornsungnoen, 2004; Almodares and Mostafafi, 2006; Almodares et al., 2006; Almodares et al., 2008d; Fazaeli et al., 2006) Sweet sorghum is harvested

Table 2. Mean comparisons among 36 sweet sorghum cultivars, lines and hybrids regarding stem yield, Brix, Sucrose and purity at university of Isfahan, Iran (Almodares and Sepahi, 1996).

Genotypes	Stem Yield (t ha⁻¹)	Brix (%)	Sucrose (%)	Purity (%)
Cultivars				
Roce	39.14	21.96	14.39	66.71
Vespa	84.53	20.99	13.05	74.59
Brandes	77.14	18.72	8.92	46.39
MN1500	83.71	20.71	12.00	57.59
E36-1	48.00	18.26	13.41	76.02
Soave	61.57	20.73	13.46	65.00
M81-E	103.57	16.01	10.26	65.10
Sumac	44.43	21.12	12.85	60.10
Sofrah	85.57	19.63	12.61	64.05
SSV-108	62.85	22.25	13.97	62.26
SSV-94	70.14	20.64	11.75	57.12
SSV-96	62.00	22.54	13.71	60.10
Theis	100.14	19.10	7.26	37.59
Foralco	97.71	20.40	12.64	60.83
Rio	95.00	22.36	16.06	71.31
S-35	58.43	19.78	11.58	58.75
Turno	39.86	11.16	6.00	35.86
Satiro	27.86	17.16	10.33	60.02
Wary	126.42	15.84	7.85	49.40
Lines				
IS 686	61.43	16.54	9.00	54.39
IS 16054	51.85	21.07	11.73	55.83
IS 18154	42.14	19.04	12.71	66.71
IS 6962	43.00	23.01	13.61	58.85
IS 9639	54.00	21.77	14.31	65.23
IS 2325	59.57	20.70	14.28	60.18
IS 6973	33.43	22.85	14.21	61.88
IS 4546	56.43	22.03	13.05	60.12
IS 19273	46.28	20.29	15.04	73.69
IS 4354	33.86	17.66	9.80	55.28
Hybrids				
A1 x IS 6973	83.28	16.46	9.53	57.17
A13 x IS 1273	97.00	21.18	14.26	66.78
A1 x IS 19261	88.13	18.69	11.82	63.04
A1 x IS 14446	87.13	16.51	10.51	62.89
A45 x IS 14446	124.13	17.95	13.36	74.06
A1 x IS 19273	128.85	17.82	13.00	73.51
A13 x IS 14446	113.56	14.32	10.73	74.40




Figure 1. Proposed layout for ethanol production and by-product from sweet sorghum.

at milk stage (Ranjbar and Almodares, 2002; Almodares et al., 2007b). Experiment crop can with-stand periods of drought better than most potential for sweet sorghum production can be increased through selection and development of adapted cultivars (Almodares and Sepahi, 1996). The aim of agronomy in sweet sorghum is to increase productivity with focus on biofuel and improved feedstock supply duration as follows.

- i) Water and fertilizer (macro- and micronutrients) effects and their interaction on sugar, grain and bagasse yield and quality.
- ii) Effect of day length, temperature and their interaction, on sugar, grain and bagasse yield and quality (and help identify suitable cultivars for season/location).
- iii) Crop rotation experiments to identify the most productive and sustainable cropping systems for different ecosystems.

Sorghum carbohydrate

The chief nonstructural carbohydrate in grain (Somani et al., 1995) of sweet sorghum is starch while sucrose is the main carbohydrate content in the stalk which is dominant form transported in the plant. Subramanian et al. (1994) have reported that cultivars with white or pale yellow seeds are most suitable for starch production. Both alpha and beta amylases (Reisi and Almodares, 2008) are sugars (Beta et al., 1995). The primary sugars

present in grain of sweet sorghum are fructose, glucose, raffinose, sucrose and maltose. In sorghum leaves, sucrose is translocated and transformed into starch during the development of grains (Smith and Frederiksen, 2000).

Grain plus stem of sweet sorghum has been shown to yield more fermentable carbohydrates than other fuel crops (Murray et al., 2008b).

In addition, the grain can be used for production of high fructose syrup (Hosseini et al., 2003) and animal feed (Ebadi et al., 1997; Azarfa et al., 1998). Therefore, sorghum is an excellent crop for biomass production. The high nonstructural carbohydrate content of its vegetative biomass can be fermented to methane or ethanol (Reddy et al., 2005). Ethanol production by fermentation of sugar solutions obtained from sweet sorghum varies widely among years at different locations, fertility (Almodares et al., 2006, 2008d), moisture, planting/harvest dates (Almodares et al., 1997a; Almodares and Mostafafi, 2006), preclude a strict linear association between number of frost free days (Almodares et al., 2007b). In stems the extent of sucrose accumulation varies among cultivars (Table 2) (Almodares and Sepahi, 1996; Almodares et al., 1997a).

Sorghum nonstructural carbohydrates contents are affected by temperature, time of day (Almodares et al., 2000), maturity (Almodares et al., 1994b), cultivar (Almodares and Sepahi, 1996), culm section, spacing and fertilization (Almodares et al., 2008d). Shading significantly reduces panicle and leaf laminae dry weights

of sorghum (Kiniry et al., 1992). Environmental conditions such as water quality (Almodares and Sharif, 2005, 2007) and also growth stage (Almodares et al., 2007c) and maturity is a factor affecting carbohydrate content (Almodares et al., 1994b). In the sweet sorghum, sucrose, glucose and fructose contents increase after anthesis (Almodares et al., 2008c). In stems, non-structural carbohydrates contents increase after preboot (Almodares et al., 2008c) and reach a maximum level near post anthesis (Almodares et al., 2008c).

Senescence and nonsenescence affect levels of sugar accumulation in the culm of sorghum cultivars (Viotor et al., 1990). The nonsenescent cultivars contain more carbohydrates at all maturity stages than the senescent cultivars. They reported that harvested senescent and nonsenescent sorghum hybrids at three stages of maturity: grain filling, black layer and post-black layer. Plants were shown to be physiologically mature when the black layer formed in the pedicel of kernels at the panicle base. McBee and Miller (1990) reported that closer spacing significantly increased total carbohydrates at the anthesis stage. Sugar production of sweet sorghum was compared with sugarcane and sugar beet and the results showed that sugar production from sweet sorghum is cheaper than both sugar cane and sugar beet (Blas et al., 2000). Therefore, it can be used as a supplementary sugar crop (Kualarni et al., 1995). So, it seems that through cultural practices, breeding and physiological manipulation can increase the carbohydrate contents in sweet sorghum plants and because of sweet sorghum has high amount of sucrose, glucose and fructose which is easily converted to ethanol by microorganism of *Saccharomyces cerevisiae*. Thus, sweet sorghum is a suitable plant for biofuel production.

The important of ethanol in biofuel

One method to reduce air pollution is to oxygenated fuel for vehicles. MTBE (Methyl tert-butyl ether) is a member of a group of chemicals commonly known as fuel oxygenates (Fischer et al., 2005). It is a fuel additive to raise the octane number. But it is very soluble in water and it is a possible human carcinogenic (Belpoggi et al., 1995). Thereby, it should be substituted for other oxygenated substances to increase the octane number of the fuel. Presently, ethanol as an oxygenous biomass fuel is considered as a predominant alternative to MTBE for its biodegradable, low toxicity, persistence and regenerative characteristic (Cassada et al., 2000). The United States gasoline supply is an ethanol blend and the importance of ethanol use is expected to increase as more health issues are related to air quality. Ethanol may be produced from many high energy crops such as sweet sorghum (Figures 1 and 2), corn, wheat, barely, sugar cane, sugar beet, cassava, sweet potato and etc (Drapcho et al., 2008). Like most biofuel crops,

sweet sorghum has the potential to reduce carbon emissions. In addition, among the plants, sweet sorghum has the following characteristics (Almodares et al., 2008e):

- i) It is an efficient converter of solar energy, as it requires low inputs and yet, a high carbohydrate producer.
- ii) As a drought-tolerant crop with multiple uses.
- iii) It has a concentration of sugar which normally varies between 12 - 21%, directly fermentable (that is, no starch to convert).
- iv) It can be cultivated in temperate, subtropical and tropical climates.
- v) All components of the plant have economic value - the grain from sweet sorghum can be used as food or feed, the leaves for forage, the stalk (along with the grain) for fuel, the fiber (cellulose) either as mulch or animal feed and with second generation technologies even for fuel.
- vi) Its bagasse, after sugar extraction, has a higher biological value than the bagasse from sugarcane, when used as feed for animals.
- vii) Its growing period is shorter (3 - 5 months) than that of sugarcane (10 - 12 months), and the quantity of water required is 1/3 of sugarcane.
- viii) In tropical irrigated areas sweet sorghum can be harvested twice each year (by ratooning) and its production can be completely mechanized.
- ix) It has some tolerance to salinity.
- x) It can produce large quantities of both readily fermentable carbohydrate and fiber per unit land area.

Therefore, based on the above characteristics, it seems that sweet sorghum is the most suitable plant for biofuel production than other crops under hot and dry climatic conditions. In addition, possible use of bagasse as a by-product of sweet sorghum include: burning to provide heat energy, paper or fiber board manufacturing, silage for animal feed or fiber for ethanol production. However, since sweet sorghum is at a relatively early stage of its development, continued research was needed to obtain better genetic material and match local agro-economic conditions. The challenge is to harvest the crop, separate it into juice and fiber and utilize each constituent for year-round production of ethanol.

Sweet sorghum juice is assumed to be converted to ethanol at 85% theoretical, or 54.4 L ethanol per 100 kg fresh stalk yield. Potential ethanol yield from the fiber is more difficult to predict (Rains et al., 1993). The emerging enzymatic hydrolysis technology has not been proven on a commercial scale (Taherzadeh and Karimi, 2008). One ton of corn grain produces 387 L of 182 proof alcohol while the same amount of sorghum grain produces 372 L (Smith and Frederiksen, 2000). Sorghum is used extensively for alcohol production (Kundiya, 1996; Bulawayo et al., 1996; Smith and Frederiksen, 2000; Gnansounou et al., 2005), where it




Figure 2. Mass balance of sweet sorghum juice extraction and ethanol production (Prasad et al., 2007).

is significantly lower in price than corn or wheat (Smith and Frederiksen, 2000). The commercial technology required to ferment sweet sorghum biomass into alcohol has been reported in china (Gnansounou et al., 2005). One ton of sweet sorghum stalks has the potential to yield 74 L of 200- proof alcohol (Smith and Frederiksen, 2000). Therefore, it seems that because ethanol can be produced from both stalk and grain of sweet sorghum (Figure 1), so it is the most suitable crop for ethanol production using for biofuel comparing to other crops such as corn or sugarcane.

Sweet sorghum stalk processing for ethanol production

Juice extraction

Juice is extracted by series of mills (Almodares et al., 2008e). The juice coming out of milling section is first screened, sterilized by heating up to 100°C and then clarified (Quintero et al., 2008). The muddy juice is then

sent to rotary vacuum filter and the filtrate juice is sent to evaporation section for concentration (syrup to ethanol). The juice can also be directly sent to fermentation section (Figure 1) (juice to ethanol). Depending on the scheme selected the juice can be concentrated using evaporators to attend various brix. In case of juice to ethanol (no syrup), it is advisable to partially increase the concentration of juice to 16 - 18 brix. The syrup which needs storage for using during off season needs to concentrate to minimum 65 brix (normally 85 brix).

Fermentation

Fermentation is a multidisciplinary process based on the chemistry, biochemistry and microbiology of the raw materials. Juice or syrup is converted into ethanol by the yeast *Saccharomyces cerevisiae*. Sugar is converted to ethanol, carbon dioxide and yeast biomass as well as much smaller quantities of minor end products such as glycerol, fusel oils, aldehydes and ketones

(Jacques et al., 1999).

Distillation and dehydration

In the distillation section, alcohol from fermented mash is concentrated up to 95% v/v. This is further concentrated to produce ethanol with 99.6% v/v (minimum) concentration. The treatment of vinasse generated in the distillation section can be done using following option:

Concentration of part of vinasse to 20 to 25% solids followed by composting using press mud available and concentration of rest of the vinasse to 55% solids and use as liquid fertilizer.

Since Iran has dry and hot climatic conditions (Almodares, 2000) therefore sweet sorghum has emerged as a leading candidate for liquid sugar (Almodares and Sepahi, 1997) and biofuel production (Nahvi et al., 1994a) with minimum inputs (Almodares and Sepahi, 1997).

Processing ethanol production from sweet sorghum grain

The ethanol production processing from sweet sorghum grain (Figure 1) is similar to corn and it can be described according to Quintero et al. (2008). After washing, crushing and milling the sweet sorghum grains, the starchy material is gelatinized, liquefied and saccharified using α -amylase and glucoamylase enzymes to produce glucose. Fermentation, distillation and dehydration processing of grain sorghum are similar to the sweet sorghum stalk. However, the by-products of grain is not similar to the stalk because DDGS (dried distillers grains with solubles) as a co-product of the ethanol production process from grain is a high nutrient feed valued which is used by the livestock industry.

Conclusion

It is clear that fuel ethanol from sweet sorghum is the best choice to be implement under hot and dry climatic conditions regarding both economic and environmental considerations. Because, sweet sorghum has higher tolerance to drought (Tesso et al., 2005), water logging and salt (Almodares et al., 2008, 2008a), alkali and aluminum soils; It may be harvested 3 - 4 months after planting (Table 1) and planted 1 - 2 times a year (in tropical areas); Its energy output / fossil energy input is higher than sugarcane, sugar beet, corn, wheat and etc... specially in temperate areas; It is more water use efficient (1/3 of water used by sugarcane at equal sugar production); Its production can be completely mecha-

nized and Its bagasse has higher nutritional value than the bagasse from sugarcane, when used for animal feeding.

Also, by implementing agricultural practices such as adequate water and fertilizers, suitable cultivars or hybrids, crop rotation, pest management and etc... can increase productivity with focus on biofuel production (Reddy et al., 2005). In addition, sweet sorghum has high amount of sucrose (Almodares and Sepahi, 1996) and invert sugar (Almodares et al., 2008c) which are easily converted to ethanol (Jacques et al., 1999; Prasad et al., 2007). Therefore, it seems that sweet sorghum is the most suitable crop for biofuel production in arid regions of the world. This awareness should push government of the countries with such climatic conditions to promote the development of projects for fuel ethanol production from sweet sorghum.

However, social aspects (including environmental concerns) should play a more significant role in the selection of the most suitable feedstocks for the alcohol industry. In this way, financial indicators would not be necessarily the decisive factors when new large-impact projects for biofuels production are studied and implemented in developing countries.

REFERENCES

- Almodares A (2000). Sweet sorghum: An energy crop for dry and hot countries. In: Proceeding of the First World Conference and Exhibition on Biomass for Energy and Industry (Ed. A Almodares). Sevilla, Spain
- Almodares A, Abdy M, Somani RB, Jilani SK (1999). Comparative study of sorghum sudangrass hybrids and lines for forage. *Ann. Plant Physiol.* 13: 6-10
- Almodares A, Aghamiri A, Sepahi A (1996). Effects of the amount and time of nitrogen fertilization on carbohydrate contents of three sweet sorghum cultivars. *Ann. Plant Physiol.* 10: 56-60
- Almodares A, Hadi MR, Ahmadpour H (2008). Sorghum stem yield and soluble carbohydrates under phenological stages and salinity levels *Afr. J. Biotech.* 7: 4051-4055
- Almodares A, Hadi MR, Dosti B (2007a). Effects of Salt Stress on Germination Percentage and Seedling Growth in Sweet Sorghum Cultivars. *J. Biol. Sci.* 7: 1492-1495
- Almodares A, Hadi MR, Dosti B (2008a). The effects of salt stress on growth parameters and carbohydrates contents in sweet sorghum. *Res. J. Environ. Sci.* 2: 298-304
- Almodares A, Hadi MR, Ranjbar M, Taheri R (2007b). The Effects of Nitrogen Treatments, Cultivars and Harvest Stages on Stalk Yield and Sugar Content in Sweet Sorghum. *Asian J. Plant Sci.* 6: 423-426
- Almodares A, Mostafafi DSM (2006). Effects of planting date and time of nitrogen application on yield and sugar content of sweet sorghum. *J. Environ. Biol.* 27: 601-605
- Almodares A, Taheri R, Adeli S (2008b). Categorization of sweet sorghum cultivars and lines as sweet, dual purpose and grain sorghum *J. Tropical. Agri.* 46: 62-63.
- Almodares A, Taheri R, Adeli S (2008c). Stalk yield and carbohydrate composition of sweet sorghum [*Sorghum bicolor* (L.) Moench] cultivars and lines at different growth stages. *J. Malesian Appl. Biol.* 37: 31-36.
- Almodares A, Sepahi A (1996). Comparison among sweet sorghum cultivars, lines and hybrids for sugar production. *Ann. Plant Physiol.* 10: 50-55
- Almodares A, Sepahi A (1997). Potential of sweet sorghum for liquid sugar production in Iran. In: Proceedings of the First International

- Sweet Sorghum Conference (Eds. A Almodares & A Sepahi). Beijing, China
- Almodares A, Sepahi A, Dalilatajary H, Gavami R (1994a). Effect of phenological stages on biomass and carbohydrate contents of sweet sorghum cultivars. *Ann. Plant Physiol.* 8: 42-48.
- Almodares A, Sepahi A, Karve A (1994b). Effect of planting date on yield and sugar production of sweet sorghum. *Ann. Plant Physiol.* 8: 49-54
- Almodares A, Sepahi A, Rezaie A (2000). Effect of breaking night period on sugar production in sweet sorghum plant. *Ann. Plant Physiol.* 14: 21-25
- Almodares A, Sepahi A, Shirvani M (1997a). The effects of planting date and genotype on carbohydrate production from sweet sorghum in the south of Iran. *Ann. Plant Physiol.* 11: 1-5
- Almodares A, Sepahi A, Shirvani M (1997b). Sweet sorghum cultural practices in Iran. In: Proceedings of the First International Sweet Sorghum Conference (Eds. A Almodares, A Sepahi & M Shirvani). Beijing, China
- Almodares A, Sharif ME (2005). Effect of water quality on yield of sugar beet and sweet sorghum. *J. Environ. Biol.* 26: 487-493
- Almodares A, Sharif ME (2007). Effects of irrigation water qualities on biomass and sugar contents of sugar beet and sweet sorghum cultivars. *J. Environ. Biol.* 28: 213-218
- Almodares A, Taheri R, Adeli S (2007c). Inter-relationship between growth analysis and carbohydrate contents of sweet sorghum cultivars and lines. *J. Environ. Biol.* 28: 527-531.
- Almodares A, Taheri R, Chung M, Fathi M (2008d). The effect of nitrogen and potassium fertilizers on growth parameters and carbohydrate content of sweet sorghum cultivars. *J. Environ. Biol.* 29: 849-852
- Almodares A, Taheri R, Hadi MR, Fathi M (2006). The Effect of Nitrogen and Potassium Fertilizers on the Growth Parameters and the Yield Components of Two Sweet Sorghum Cultivars. *Pakistan. Biol. Sci.* 9: 2350-2353
- Almodares A, Taheri R, Safavi V (2008e). Sorghum, Botany, Agronomy and Biotechnology, Jahad Daneshgahi of University of Isfahan, Isfahan
- Anglani C (1998) Sorghum carbohydrates - A Rev. *Plant Food Human Nutr.* 52: 77-83
- Azarfa A, Ghorbani G, Asadi Y, Almodares A (1998). Dry matter, organic matter and starch degradability in barley and three low, medium and high tannin sorghum varieties. *Agric. Sci.* 8: 56-60
- Belpoggi F, Soffritti M, Maltoni C (1995). Methyl tertiary-butyl ether (MTBE) - A gasoline additive - causes testicular and lymphohematopoietic cancers in rats. *Toxicol. Ind. Health* 11: 119-149.
- Beta T, Rooney L, Waniska R (1995). Malting characteristics of sorghum cultivars. *Cereal Chem.* 72: 533-538
- Blas J, Almodares A, Somani RB (2000). Econo- Feasibility of sugar production from sweet sorghum. *PKV Res. J.* 24: 30-34
- Bulawayo B, Bvochora JM, Muzondo MI, Zvauya R (1996). Ethanol production by fermentation of sweet sorghum juice using various yeast strains. *World J. Microb. Biot.* 12: 357-360.
- Cassada DA, Zhang Y, Snow DD, Spalding RF (2000). Trace analysis of ethanol, MTBE, and related oxygenate compounds in water using solid-phase microextraction and gas chromatography/mass spectrometry. *Anal. Chem.* 72: 4654-4658.
- Drapcho CM, Nhuan NP, Walker TH (2008). *Biofuels Engineering Process Technol.*, The McGraw-Hill companies, Inc, USA
- Ebadi M, Reza RP, Khorvash M, Adl KN, Almodares A (1997). Nutrient composition and metabolizable energy values of 36 sorghum grain cultivars and two types of corn 1: 67-76
- Fazaeli H, Golmohammadi HA, Almodares A, Mosharraf S, Shaei A (2006). Comparing the performance of sorghum silage with maize silage in feedlot calves. *Pakistan J. Biol. Sci.* 9: 2450-2455
- Fischer A, Oehm C, Selle M, Werner P (2005). Biotic and abiotic transformations of methyl tertiary butyl ether (MTBE). *Environ. Sci. Pollut. Res. Int.* 12: 381-386
- Gnansounou E, Dauriat A, Wyman CE (2005). Refining sweet sorghum to ethanol and sugar: economic trade-offs in the context of North China. *Bioresource Technol.* 96: 985-1002
- Hosseini H, Almodares A, Miroliaei M (2003). Production of fructose from sorghum grain. In: Proceeding of the 11th Iranian Biology Conference (Eds. H Hosseini, A Almodares & M Miroliaei). Urmia, Iran
- Jacques K, Lyons TP, Kelsall DR (1999). *The Alcohol Textbook*. 3rd Eds. p 388
- Jafarinia M, Almodares A, Khorvash M (2005). Using sweet sorghum bagasse in silo In: Proceeding of the 2nd Congress of Using Renewable Sources and Agric. Wastes (Eds. M Jafarinia, A Almodares & M Khorvash). Khorasgan Azade University, Isfahan, Iran
- Kiniry JR, Tischler CR, Rosenthal WD, Gerik TJ (1992). Nonstructural carbohydrate utilization by sorghum and maize shaded during grain growth. *Crop Sci.* 32: 131-137.
- Kualarni DP, Almodares A, Somani RB (1995). Sweet sorghum- A supplementary sugar crop in Iran. *Ann. Plant Physiol.* 9: 90-94
- Kundiayana DK (1996). Sorghanol: In-field production of ethanol from sweet sorghum. In: Dairy Sci. College. University of Agricultural Science Bangalore, India
- McBee G, Miller FR (1990). Carbohydrate and lignin partitioning in sorghum stems and blades. *Agron. J.* 82: 687-690.
- Murray SC, Rooney WL, Mitchell SE, Sharma A, Klein PE, Mullet JE, Kresovich S (2008a). Genetic Improvement of Sorghum as a Biofuel Feedstock: II. QTL for Stem and Leaf Structural Carbohydrates. *Crop Sci.* 48: 2180-2193
- Murray SC, Sharma A, Rooney WL, Klein PE, Mullet JE, Mitchell SE, Kresovich S (2008b). Genetic Improvement of Sorghum as a Biofuel Feedstock: I. QTL for Stem Sugar and Grain Nonstructural Carbohydrates. *Crop Sci.* 48: 2165-2179
- Nahvi I, Almodares A, Motocalemi M (1994a). Comparing ethanol production from sweet sorghum juice and sugar beet molasses. In: Proceeding of the 1st Iranian Congress on Chemical Engineering (Eds. I Nahvi, A Almodares & M Motocalemi). Tehran, Iran
- Nahvi I, Almodares A, Rasuli A (1994b). The effect of environ. factors on ethanol production from sweet sorghum juice. In: Proceeding of the 11th Iranian Congress on Chem. Eng. (Eds. I Nahvi, A Almodares & A Rasuli). Tehran, Iran
- Pholsen S, Sornsungnoen N (2004). Effects of nitrogen and potassium rates and planting distances on growth, yield and fodder quality of forage sorghum (*Sorghum bicolor* L. Moench). *Pakistan J. Biol. Sci.* 7: 1793-1800
- Prasad S, Singh A, Jain N, Joshi HC (2007). Ethanol production from sweet sorghum syrup for utilization as automotive fuel in India. *Energy Fuels* 21: 2415-2420.
- Quintero JA, Montoya MI, Sanchez OJ, Giraldo OH, Cardona CA (2008). Fuel ethanol production from sugarcane and corn: Comparative analysis for a Colombian case. *Energy* 33: 385-399
- Rains GC, Cundiff JS, Welbaum GE (1993). Sweet sorghum for a piedmont ethanol industry. In: *New crops* (Eds. J Janick & JE Simon). Wiley New York
- Ranjbar M, Almodares A (2002). The effect of nitrogen application time and harvesting on sweet sorghum yield. *Flavarjan Azad University* 1: 59-69
- Reddy BVS, Ramesh S, Reddy PS, Ramaiah B, Salimath PM, Kachapur R (2005). Sweet Sorghum—A Potential Alternate Raw Material for Bio-ethanol and Bio-energy. *Int. Sorghum Millets Newslett.* 46: 79–86
- Rego TJ, Rao VN, Seeling B, Pardhasaradhi G, Rao JVDK (2003). Nutrient balances a guide to improving sorghum and ground based dry land cropping systems in semi-arid topical India. *Field Crop Res.* 81: 53-68.
- Reisi F, Almodares A (2008). The effect of planting date on amylose content in sorghum and corn. In: Proceeding of the 3rd Int. Biol. conference (Eds. F Reisi & A Almodares). Tehran, Iran
- Rezaie A, Almodares A, Amini M, Khanghly S (2005). The effect of milo photoperiodic to red light In: Proceeding of the 1st National Congress on Forage Grass (Eds. A Rezaie, A Almodares, M Amini & S Khanghly). University of Tehran Agricultural and Natural Research, Karaj, Iran
- Smith CW, Frederiksen RA (2000). *Sorghum: Origin, history, technology, and production*, John Wiley and Sons, New York.
- Somani RB, Almodares A, Shirvani M (1995). Preliminary studies on sweetener production from sorghum grains. *Ann. Plant Physiol.* 9: 146-148
- Subramanian V, Hosney RC, Cox PB (1994). Factors affecting the

- color and appearance of sorghum starch. *Cereal Chem.* 71: 275-278
- Taherzadeh MJ, Karimi K (2008). Bioethanol: Market and Production Processes. In: *Biofuels Refining and Performance* (Ed. A Nag). The Mc Graw- Hill, USA
- Tesso TT, Clafin LE, Tuinstra MR (2005). Analysis of Stalk Rot Resistance and Genetic Diversity among Drought Tolerant Sorghum Genotypes. *Crop Sci.* 45: 645-652
- Tsialtas JT, Maslaris N (2005). Effect of N Fertilization Rate on Sugar Yield and Non-Sugar Impurities of Sugar Beets (*Beta vulgaris*) Grown Under Mediterranean Conditions. *J. Agron. Crop Sci.* 191: 330-339
- Vietor DM, Miller FR, Cralle HT (1990). Nonstructural carbohydrates in axillary branches and main stem of senescent and nonsenescent sorghum types. *Crop Sci.* 30: 97-100